

Alive?

- Check pulse and respirations
- Stay with your patient.

get Loud!

- PUSH the code button.
- SHOUT OUT LOUD for help.

Initiate CPR

- Start immediately and minimize interruptions.
- Push hard and fast.
- Use the backboard when it arrives.

Voltage

- Turn on AED/Defibrillator.
- Apply pads.
- Shock if advised.

Epi

- Confirm intravenous access.
- Open the crash cart.
- Prepare to give pre-filled epinephrine syringe.

ALIVE?

get LOUD

Initiate CPR

Voltage

Epi



Alive?

- Check pulse and respirations
- Stay with your patient.




get Loud

- PUSH the code button.
- SHOUT OUT LOUD for help.



Initiate CPR

- Start immediately and minimize interruptions.
- Push hard and fast.
- Use the backboard when it arrives.



Voltage

- Turn on AED/Defibrillator.
- Apply pads.
- Shock if advised.



Epinephrine

- Confirm intravenous access.
- Open the crash cart.
- Prepare to give pre-filled epinephrine syringe.

Content

Assumptions – starts w/ unresponsive patient, no one is ACLS certified; no APP/MD present.

- Scripting for Pulse check may be unreliable – when in doubt get Loud and Initiate CPR.
- Scripting for SHOUT OUT LOUD – Patient coding. Bring the crash cart. I need help. (No need to add please and thank you).
- Remind – stayin alive 100-120/minute; minimize interruptions; choreograph the backboard– using the crash cart; bed mechanics/move the rails/head of bed positioning; choreograph the
- Voltage – defib vs. aed– if in AED mode, follow directions; If ACLS – open door and do your thing; pad placement to not mess with CPR; keeping CPR going while charging (hold for analyze only); Say -- Verbal clear– I'm clear, you're clear, we are all clear AND Visual clear ; Say -- Shock Delivered. Restart CPR now.
- EPI – confirm IV access & be ready to communicate; OPEN THE CRASH CART AND FIND EPI in the 2nd drawer on the left side – beige/ecru box; open from bottom where it says open, placing your thumb in the `/1 moon. Flip/Flip/Twist/Twist. If no acls, please assemble the epi and be prepared to give when ACLS provider instructs if you are credentialed to give iv meds. If ACLS certified personnel present in the room– give 1mg 1:10,000 epi IV PUSH FAST(THE WHOLE THING/abbject– 10ml-- FOR YOUR ADULT PATIENT...) if you are credentialed to give iv medications.
- General stuff – closed-loop communication. Roles ... please. Some of these tasks should all be happening simultaneously for example when there are enough people-- Once CPR starts never stop. Voltage and epi.